

RECORDS OF THE GLOBALLY THREATENED RUSTY-SPOTTED CAT IN ODISHA, INDIA

Himanshu S. Palei^{1*}, Nimain C. Palei², Bhakta P. Rath², Arun K. Mishra³

¹North Orissa University, India

²Office of the PCCF (Wildlife) and Chief Wildlife Warden, India

³Sundargarh Forest Division, India

*e-mail: himanshu.palei@gmail.com

Received: 22.04.2019. Revised: 12.07.2019. Accepted: 13.07.2019.

The Rusty-spotted Cat, *Prionailurus rubiginosus*, is a vulnerable species, endemic to India, Nepal and Sri Lanka. The goal of the study is to provide an overview of the current distribution of the Rusty-spotted Cat in Odisha state through camera trap survey, review of published scientific literature and rescue records. This study presents 14 localities including seven new localities from the Odisha state in India. A long-term study, habitat protection and studying its ecology are recommended for initiating further steps to conserve its range.

Key words: camera traps, conservation, distribution range, *Prionailurus rubiginosus*, small cats, threats

Introduction

The Rusty-spotted Cat *Prionailurus rubiginosus* (I. Geoffroy Saint-Hilaire, 1831) is the smallest feline in the world and endemic to India, Nepal and Sri Lanka (Mukherjee et al., 2016). It has been predicted that the global population may decline by 20–25% in the coming decades corresponding to a loss and degradation of the remaining habitat (Mukherjee et al., 2016). The preferred habitat for this species is associated with dense vegetation in moist and dry deciduous forest as well as scrub and grassland (Patel, 2011). Further, there are also reports of its occurrence within agricultural areas and human settlements (Nowell & Jackson, 1996; Mukherjee, 1998; Nekaris, 2003; Athreya, 2010). The Rusty-spotted Cat's status, distribution and ecology are still unknown because of its elusive behaviour and naturally low density (Mukherjee et al., 2016).

This species is affected by habitat loss and 75% of its habitat in the current distributional range facing an imminent danger of conversion of forest to agriculture and industry (Mukherjee et al., 2016). According to the International Union for Conservation of Nature (IUCN), the Rusty-spotted Cat is now listed in the «Near Threatened» category because of its larger range than previously known. It is protected under the Schedule I of the Indian Wildlife (Protection) Act, 1972. However, a lack of systematic surveys on its status has designated it as one of the lesser known species throughout its distributional range (Mukherjee et al., 2016).

In India, this species is found in an irregular manner from the extreme south of Tamil Nadu to the extreme north of Jammu and Kashmir and from the extreme west of Gujarat to Odisha (formerly known as Orissa) in the east, through Madhya Pradesh in the central region (Patel & Jackson, 2005; Mali & Srinivasulu, 2015; Nayak et al., 2017). Although the Rusty-spotted Cat's distribution covers the large part of Odisha, eastern India, there are few records in this state (Palei & Debata, 2019). This may be due to the ecological feature of the species, such as naturally low density and secretive behaviour (Mukherjee et al., 2016), but probably reflect the lack targeted surveys by biologists which is evident from the recent faunal inventories (Mohapatra et al., 2014; Debata et al., 2013; Debata et al., 2015; Palei H. et al., 2018). Here we present a review of the occurrence and current distribution range of the Rusty-spotted Cat and also add some distribution records from Odisha, eastern India.

Material and Methods

Odisha State is located between 17.49–22.34°N and 81.29–87.29°E covering an area of 155 707 km² along the eastern coast of India (Fig. 1). Odisha falls under the Deccan Peninsula biogeographic zone spreading over the Deccan plateau, the Central highlands, the Eastern Ghats, the Gangetic plain and Coasts biogeographic province (Sinha, 1971). Most precipitation falls during the monsoon, with a mean annual total rainfall of 1451 mm. Temperatures range from a minimum of 2°C in winter to a maximum of 45°C in summer.




Fig. 1. Map showing the locations of the Rusty-spotted Cat *Prionailurus rubiginosus* occurrence in Odisha, eastern India.

To assess the current distribution status of the Rusty-spotted Cat in Odisha, we reviewed available literature including published articles, books and unpublished reports. We also gathered records from forest department rescues and camera trap surveys conducted in different part of Odisha. Camera traps were deployed in forest patches in seven forest divisions and seven wildlife sanctuaries (Table 1). The study was carried out for terrestrial mammals from January 2017 to December 2018 in Odisha (Palei N. et al., 2018). Camera traps were installed at 493 sampling stations in forest plains to hilly terrain along the animal trails, nullahs as well as rocky and sandy stream beds (Table 1). Each camera trap station was composed of two camera traps (Cuddeback Model C1). All camera traps were strapped to trees approximately 40 cm above ground. Camera traps were set to operate 24 h per day and programmed to delay sequential photographs by 30 s recording time. Each camera trap was checked at least once a week for battery level, positioning and to replace memory (SD) cards. Each photograph was manually checked to identify the species. Date, time and temperature were noted for each identified species.

Results and Discussion

During the survey we recorded a total of 14 localities, including seven published records, two rescue records and five camera trap records (Table 2, Fig. 1, Fig. 2). Out of these 14 localities, seven localities were in Protected Areas (Fig. 1). The Rusty-spotted Cat was previously known from seven localities. So, in this study, we recorded seven more localities of this species from different parts of Odisha (Fig. 2). All these new localities were recorded between 2016 and 2018. The total sampling had 493 camera-trap stations, with a total sampling effort of 12 325 trap days, considering all camera-trap stations (Table 1). We obtained five Rusty-spotted Cat photos in different camera trap stations (Table 1, Table 2).

As mentioned in Material and Methods, camera traps were deployed to gather information on terrestrial mammals and was not species-specific. As a result, these camera traps were installed at a height of 40 cm above ground. There are possibilities of missing out small carnivore species like the Rusty-spotted Cat. The detection probability of the Rusty-spotted Cat might be different and hence there may be a difference between estimated photo captured and actual captured in Table 1.

Table 1. Summary of camera trap capture records of Rusty-spotted Cat *Prionailurus rubiginosus* in different forest divisions, Odisha, India

No.	Name of forest division	Camera trap stations	Trap days	Total sampling effort	Rusty-spotted Cat photo captured
1	Sundargarh Forest Division	45	25	1125	2
2	Debrigarh Wildlife Sanctuary	39	25	975	1
3	Baragarh Forest Division	45	25	1125	1
4	Balangir Forest Division	37	25	925	0
5	Sunabeda Wildlife Sanctuary	35	25	875	0
6	Hadagarh Wildlife Sanctuary	40	25	1000	1
7	Kuldiha Wildlife Sanctuary	40	25	1000	0
8	Karlapat Wildlife Sanctuary	34	25	850	0
9	Kalahandi North Forest Division	21	25	525	0
10	Rourkela Forest Division	33	25	825	0
11	Bonai Forest Division	35	25	875	0
12	Khariar Forest Division	38	25	950	0
13	Khalasuni Wildlife Sanctuary	26	25	650	0
14	Badrama Wildlife Sanctuary	25	25	625	0
Total		493	25	12325	5



Fig. 2. Occurrence records of the Rusty-spotted Cat *Prionailurus rubiginosus* in Odisha, eastern India. **A.** Rescued in Ghumusur North Forest Division; **B.** Rescued in Phulbani Forest Division; **C.** Camera trapped in Debrigarh Wildlife Sanctuary; **D.** Camera trapped in Bargarh Forest Division; **E.** Camera trapped in Sundargarh Forest Division; **F.** Camera trapped in Sundargarh Forest Division; **G.** Camera trapped in Bolangir Forest Division (see details in Table 2).

Table 2. All available records on the occurrence of the Rusty-spotted Cat *Prionailurus rubiginosus* in Odisha, India (numbers of locations correspond with numbers in Fig. 1)

No.	Year	Locations	Forest type	Type of records	Sources
1	1969	Tuluka Reserve Forest near Purunakote village of Satkosia Tiger Reserve, Satkosia Wildlife Division, Angul District	Moist deciduous forest	Published records	Write, 1984
2	1995	Bhramaramadi village near Daringibadi of Baliguda Forest Division, Phulabani District	Moist deciduous forest	Published records	Acharjyo et al., 1997
3	2010	Karlapat Wildlife Sanctuary of Kalahandi South Wildlife Division, Kalahandi District	Dry deciduous forest	Published records	Palei & Debata, 2019
4	2013	Chandanpur village near Dukura range of Similipal Tiger Reserve, Baripada Forest Division, Mayurbhanj District	Moist deciduous forest	Published records	Palei & Debata, 2019
5	2014	On a forest road leading to Jhagadabehera village of Lakhanpur range, Hirakud Wildlife Division, Sambalpur District	Dry deciduous forest	Published records	Palei & Debata, 2019
6	2017	Dangadiha beat, Kaptipada range, Similipal Tiger Reserve, Baripada Forest Division, Mayurbhanj District	Moist deciduous forest	Published records	Mishra et al., 2019
7	2017	Dangadiha beat, Kaptipada range, Similipal Tiger Reserve, Baripada Forest Division, Mayurbhanj District	Moist deciduous forest	Published records	Mishra et al., 2019
8	2017	Jhagadabehera village of Lakhanpur range, Debrigarh Wildlife Sanctuary, Hirakud Wildlife Division, Sambalpur District	Dry deciduous forest	Camera trap	Present study
9	2017	Binjigiri, Bhanjanagar, Ghumusur North Forest Division, Ganjam District	Moist deciduous forest	Rescued	Present study
10	2018	Keredi, Suradei, Phulbani Forest Division, Phulbani District	Moist deciduous forest	Rescued	Present study
11	2018	Jamseth, Gandhmardhan Proposed Reserve Forest, Nrusinghanath range of Bargarh Forest Division, Bargarh District	Dry deciduous forest	Camera trap	Present study
12	2018	Dangakhoh, Garjanpahad Reserve Forest, Dagora, Hemgiri, Sundargarh Forest Division, Sundargarh District	Dry deciduous forest	Camera trap	Present study
13	2018	Telianala of Chengapahad, Kanika, Hemgiri Reserve Forest, Sundargarh Forest Division, Sundargarh District	Dry deciduous forest	Camera trap	Present study
14	2018	Jhaliati, Lathore, Bolangir Forest Division, Bolangir District	Dry deciduous forest	Camera trap	Present study

The Rusty-spotted Cat is relatively rare in comparison to other small cats found in Odisha (Palei et al., 2016; Palei H. et al., 2018; Palei & Debata, 2019). However, it is found to be widespread in Odisha. The population of the Rusty-spotted Cat is threatened from human induced activities mainly conversion of forest habitat for development. The last seven decades 40.5% of the forest cover was declined in Odisha (Reddy et al., 2013). Deforestation started mainly after the development of infrastructure, dam construction and intense population growth after country's independence (Reddy et al., 2013).

For effective conservation and management of globally threatened species, information on the population status, movement patterns, diet, habitat requirements and demography as well as population status and ecology of prey species is required (Grassman et al., 2005). Unlike to other congeners, little is known about the natural history and population status of the Rusty-spotted Cat. Similar to our study, most of the data on the Rusty-spotted Cat are available mainly from opportunistic sightings and bycatch camera trap images from large carnivore study (Mukherjee et al., 2016). No studies have been targeted on the Rusty-spotted Cat in its range. Therefore, its population status

and identification of known and possible threats for this cat are highly recommended.

The recent increase in occurrence of the Rusty-spotted Cat from different locations in Odisha and elsewhere in India is of concern and necessitates awareness raising programmes, particularly for locals and forest departments who occasionally misidentified the animal with the Jungle Cat *Felis chaus* Schreber, 1777 and leopard cub *Panthera pardus* (Linnaeus, 1758) (Palei & Debata, 2019). The lack of scientific understanding of the Rusty-spotted Cat in Odisha and elsewhere in its range restricts the ability to conserve the species. Identification and conservation of the Rusty-spotted Cat key habitats can play an important role in conservation planning for the species. Therefore, we recommend to conduct a large-scale habitat modelling study for a better understanding of its potential distribution range, not only within the Odisha state, but broader, in its total distributional range.

Acknowledgements

We are grateful to Dr. Sandeep Tripathi (Principal Chief Conservator of Forests (Wildlife) and Chief Wildlife Warden, Odisha Forest Department) for supporting and granting us permission to carry out the study. We are thankful to Divisional Forest Officers and all forest officials for supporting the study.

References

- Acharjyo L.N., Purohit K.L., Patnaik S.K. 1997. Occurrence of the Rusty-spotted Cat in Orissa. *Journal of Bombay Natural History Society* 94: 554–555.
- Athreya V. 2010. Rusty-spotted Cat more common than we think? *Cat News* 53: 27.
- Debata S., Palei H.S., Mohapatra P.P., Palita S.K. 2015. Additional Records of Cantor's Leaf-Nosed Bat *Hipposideros galeritus* Cantor, 1846 (Mammalia: Chiroptera: Hipposideridae) in Eastern India: Odisha. *Journal of Threatened Taxa* 7(8): 7477–7479. DOI: 10.11609/JoTT.o4247.7477-9
- Debata S., Palei H.S., Mohapatra P.P., Mishra A.K. 2013. First record of Lesser False Vampire Bat (*Megaderma spasma*, Linnaeus, 1758) from Sundargarh, Odisha, India. *Small Mammal Mail* 5(1): 26–27.
- Grassman Jr. L.I., Tewes M.E., Silvy N.J., Kreetiyutanont K.J. 2005. Ecology of three sympatric felids in a mixed evergreen forest in north-central Thailand. *Journal of Mammalogy* 86(1): 29–38. DOI: 10.1644/1545-1542(2005)086<0029:EOTSFI>2.0.CO;2
- Mali S., Srinivasulu C. 2015. Records of Rusty-Spotted Cat, Sri Lankamalleshwara Wildlife Sanctuary India. *Cat News* 62: 20–21.
- Mishra S.R., Mohan M., Pati J. 2019. Camera trap records of rusty-spotted cat from Similipal Tiger Reserve, Odisha, India. *Cat News* 69: 26–27.
- Mohapatra P.P., Palei H.S., Hussain S.A. 2014. Occurrence of Asian Small-Clawed Otter *Aonyx cinereus* (Illiger, 1815) in Eastern India. *Current Science* 107(3): 367–370.
- Mukherjee S. 1998. Cats: some large many small. In: S.K. Mukherjee (Ed.): *Small Cats of India. Envis Bulletin*. Vol. 1(1). Dehradun: Wildlife Institute of India. P. 5–13.
- Mukherjee S., Duckworth J.W., Silva A., Appel A., Kittle A. 2016. *Prionailurus rubiginosus*. In: *The IUCN Red List of Threatened Species 2016: e.T18149A50662471*. Available from <http://dx.doi.org/10.2305/IUCN.UK.2016-1.RLTS.T18149A50662471.en>
- Nayak S., Shah S., Borah J. 2017. First record of Rusty-spotted Cat *Prionailurus rubiginosus* (Mammalia: Carnivora: Felidae) from Ramgarh-Vishdhari Wildlife Sanctuary in semi-arid landscape of Rajasthan, India. *Journal of Threatened Taxa* 9(1): 9761–9763. DOI: 10.11609/jott.3303.9.1.9761-9763
- Nekaris K.A.I. 2003. Distribution and behaviour of three small wild cats in Sri Lanka. *Cat News* 38: 30–32.
- Nowell K., Jackson P. 1996. *Wild Cats: Status Survey and Conservation Action Plan*. Gland, Switzerland: IUCN. 383 p.
- Palei H.S., Debata S. 2019. On the Occurrence of Rusty-Spotted Cat *Prionailurus rubiginosus* (I. Geoffroy Saint-Hilaire, 1831) (Mammalia: Carnivora: Felidae) in Odisha, Eastern India. *Proceedings of the Zoological Society* 72(1): 102–105. DOI: 10.1007/s12595-017-0243-x
- Palei H.S., Das U.P., Debata S. 2018. The vulnerable fishing cat *Prionailurus viverrinus* in Odisha, eastern India: status and conservation implications. *Zoology and Ecology* 28(2): 69–74. DOI: 10.1080/21658005.2018.1468646
- Palei H.S., Pradhan T., Sahu H.K., Nayak A.K. 2016. Estimating mammalian abundance using camera traps in the tropical forest of Similipal Tiger Reserve, Odisha, India. *Proceedings of the Zoological Society* 69(2): 181–188. DOI: 10.1007/s12595-015-0143-x
- Palei N.C., Rath B.P., Palei H.S., Mishra A.K. 2018. Occurrence of melanistic leopard in Odisha, eastern India. *Cat News* 68: 7–8.
- Patel K. 2011. Preliminary survey of small cats in Eastern Gujarat, India. *Cat News* 54: 8–11.
- Patel K., Jackson P. 2005. Rusty-spotted cat in India: new distribution data. *Cat News* 42: 27.
- Reddy C.S., Jha C.S., Dadhwal V.K. 2013. Assessment and monitoring of long-term forest cover changes in Odisha, India using remote sensing and GIS. *Environmental Monitoring and Assessment* 185(5): 4399–4415. DOI: 10.1007/s10661-012-2877-5
- Sinha B.N. 1971. *Geography of Orissa*. New Delhi: National Book Trust. 172 p.
- Wright A. 1984. A note on the wild cats of the north-eastern region of India. In: P. Jackson (Ed.): *The plight of the cats*. Madhya Pradesh: IUCN/SSC Cat Specialist Group. P. 81–84.

НАХОДКИ ГЛОБАЛЬНО УГРОЖАЕМОГО ВИДА РЖАВОЙ КОШКИ В ОДИШЕ (ИНДИЯ)

Х. С. Палей^{1,*}, Н. Ч. Палей², Б. П. Пат², А. К. Мишра³

¹Университет Северной Ориссы, Индия

²Офис главного управления по сохранению лесов в Одисше, Индия

³Лесной отдел в Сундаргархе, Индия

*e-mail: himanshu.palei@gmail.com

Ржавая кошка, *Prionailurus rubiginosus*, является уязвимым видом, эндемиком для Индии, Непала и Шри Ланки. Целью исследования явилось представить обзор современного распространения ржавой кошки в штате Одисша (Индия) с использованием данных фотоловушек, литературы, и находок спасенных животных. Данное исследование представляет 14 местонахождений, включая семь новых для штата Одисша. Долговременное изучение, охрана местообитаний и экологии вида необходимы для осуществления следующих шагов по сохранению ржавой кошки на этой территории.

Ключевые слова: *Prionailurus rubiginosus*, ареал, малые кошки, сохранение, угрозы, фотоловушки